

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

This listing of claims will replace all prior versions and listings of claims in the application:

LISTING OF CLAIMS

1. (Previously presented) A compound of formula I



(I)

wherein

A is N;

X is CR₁₁ or N;

Y is CR₇ or N with the proviso that when X is N; then Y must be CR₇;

R₁ is H, C₁-C₆alkylcarbonyl, C₁-C₆alkylcarbonyloxy or an C₁-C₆alkyl, C₁-C₆alkenyl, C₁-C₆alkynyl or cycloheteroalkyl group each optionally substituted;

R₂, R₃, R₄, R₅ and R₆ are each independently H, halogen, OH or an optionally substituted C₁-C₆alkyl group;

R₇ and R₁₁ are each independently H, halogen or an C₁-C₆alkyl, aryl, heteroaryl or C₁-C₆alkoxy group each optionally substituted;

R₈ is an C₁-C₆alkyl, aryl or heteroaryl group each optionally substituted;

R₉ is H, halogen or an C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkenyl, aryl or heteroaryl group each optionally substituted;

R₁₀ is H, OH or an optionally substituted C₁-C₆alkoxy group;

m is an integer of 2;

n is 0 or an integer of 1, 2 or 3; and

--- represents a single bond or a double bond; or
a pharmaceutically acceptable salt thereof.

2. (Cancelled)

3. (Original) The compound according to claim 1 wherein R₈ is an optionally substituted phenyl group.

B' 4. (Original) The compound according to claim 1 wherein R₂, R₃, R₄, R₅ and R₆ are H.


5. (Previously presented) The compound according to claim 1 wherein R₁ is H or a C₁-C₆alkyl or cycloheteroalkyl group each optionally substituted.

6. (Original) The compound according to claim 5 selected from the group consisting of:

1-(phenylsulfonyl)-4-piperazin-1-yl-1H-indole;
1-[(2-bromophenyl)sulfonyl]-4-piperazin-1-yl-1H-indole;
1-[(6-chloroimidazo[2,1-b][1,3]thiazol-5-yl)sulfonyl]-4-piperazin-1-yl-1H-indole;
1-[(3,4-dimethoxyphenyl)sulfonyl]-4-piperazin-1-yl-1H-indole;
1-[(5-chloro-3-methyl-1-benzothien-2-yl)sulfonyl]-4-piperazin-1-yl-1H-indole;
1-[(4-bromophenyl)sulfonyl]-4-piperazin-1-yl-1H-indole;
1-[(5-bromothien-2-yl)sulfonyl]-4-piperazin-1-yl-1H-indole;
1-[(4,5-dichlorothien-2-yl)sulfonyl]-4-piperazin-1-yl-1H-indole;
methyl 4-[(4-piperazin-1-yl-1H-indol-1-yl)sulfonyl]phenyl ether;
4-piperazin-1-yl-1-[[4-(trifluoromethoxy)phenyl]sulfonyl]-1H-indole;
4-(4-benzylpiperazin-1-yl)-1-(phenylsulfonyl)-1H-indole;
4-(4-benzylpiperazin-1-yl)-1-[(2-bromophenyl)sulfonyl]-1H-indole;
4-(4-benzylpiperazin-1-yl)-1-[(6-chloroimidazo[2,1-b][1,3]thiazol-5-yl)sulfonyl]-1H-indole;
4-(4-benzylpiperazin-1-yl)-1-[(3,4-dimethoxyphenyl)sulfonyl]-1H-indole;
4-[4-(3-methoxybenzyl)piperazin-1-yl]-1-(phenylsulfonyl)-1H-indole;
1-(phenylsulfonyl)-4-[4-(pyridin-4-ylmethyl)piperazin-1-yl]-1H-indole;
1-(phenylsulfonyl)-4-[4-(pyridin-3-ylmethyl)piperazin-1-yl]-1H-indole;
1-[(2-bromophenyl)sulfonyl]-4-[4-(3-methoxybenzyl)piperazin-1-yl]-1H-indole;
1-[(2-bromophenyl)sulfonyl]-4-[4-(pyridin-4-ylmethyl)piperazin-1-yl]-1H-indole;
1-[(2-bromophenyl)sulfonyl]-4-[4-(pyridin-3-ylmethyl)piperazin-1-yl]-1H-indole;
1-(phenylsulfonyl)-5-piperazin-1-yl-1H-indazole;
1-(phenylsulfonyl)-6-piperazin-1-yl-1H-indazole;

B/ 1-[(2-bromophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
1-[(4-bromophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
1-[(4-bromophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
1-[(5-bromothien-2-yl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
1-[(5-bromothien-2-yl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
1-[(4-fluorophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
1-[(4-fluorophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
methyl 4-[(5-piperazin-1-yl-1H-indazol-1-yl)sulfonyl]phenyl ether;
1-phenylsulfonyl-4-(4-propylpiperazin-1-yl)-1H-indazole;
1-phenylsulfonyl-4-piperazin-1-yl-1H-indazole;
1-phenylsulfonyl-4-(4-phenethylpiperazin-1-yl)-1H-indazole;
1-phenylsulfonyl-4-[4-(3-phenylpropyl)-piperazin-1-yl]-1H-indazole; and
the pharmaceutically acceptable salts thereof.

7. (Currently amended) A method for the treatment of a disorder of the central nervous system related to or affected by the 5-HT₆ receptor wherein said disorder is selected from the group consisting essentially of: schizophrenia and depression; ~~and a cognitive disorder~~ in a patient in need thereof which comprises administering to said patient a therapeutically effective amount of a compound of formula I.



wherein

A is N;

X is CR₁₁ or N;

Y is CR₇ or N with the proviso that when X is N, then Y must be CR₇;

R₁ is H, C₁-C₆alkylcarbonyl, C₁-C₆alkylcarbonyloxy or an C₁-C₆alkyl, C₁-C₆alkenyl, C₁-C₆alkynyl or cycloheteroalkyl group each optionally substituted;

R_2, R_3, R_4, R_5 and R_6 are each independently H, halogen, OH or an optionally substituted C_1 - C_6 alkyl group;

R_7 and R_{11} are each independently H, halogen or an C_1 - C_6 alkyl, aryl, heteroaryl or C_1 - C_6 alkoxy group each optionally substituted;

R_8 is an C_1 - C_6 alkyl, aryl or heteroaryl group each optionally substituted;

R_9 is H, halogen or an C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkenyl, aryl or heteroaryl group each optionally substituted;

R_{10} is H, OH or an optionally substituted C_1 - C_6 alkoxy group;

m is an integer of 2;

n is 0 or an integer of 1, 2 or 3; and

--- represents a single bond or a double bond; or

a pharmaceutically acceptable salt thereof.


8. (Cancelled)

9. (Cancelled)

10. (Cancelled)

11. (Cancelled)

12. (Previously presented) A pharmaceutical composition which comprises a pharmaceutically acceptable carrier and an effective amount of a compound of formula I.



(I)

wherein

A is N;

X is CR_{11} or N;

Y is CR₇ or N with the proviso that when X is N, then Y must be CR₇;

R₁ is H, C₁-C₆alkylcarbonyl, C₁-C₆alkylcarbonyloxy or an C₁-C₆alkyl, C₁-C₆alkenyl, C₁-C₆alkynyl or cycloheteroalkyl group each optionally substituted;

B/ R₂, R₃, R₄, R₅ and R₆ are each independently H, halogen, OH or an optionally substituted C₁-C₆alkyl group;

R₇ and R₁₁ are each independently H, halogen or an C₁-C₆alkyl, aryl, heteroaryl or C₁-C₆alkoxy group each optionally substituted;

R₈ is an C₁-C₆alkyl, aryl or heteroaryl group each optionally substituted;

R₉ is H, halogen or an C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkenyl, aryl or heteroaryl group each optionally substituted;

R₁₀ is H, OH or an optionally substituted C₁-C₆alkoxy group;

m is an integer of 2;

n is O or an integer of 1, 2 or 3; and

--- represents a single bond or a double bond; or

a pharmaceutically acceptable salt thereof.

13. (Cancelled)

14. (Original) The composition according to claim 12 wherein R₈ is an optionally substituted phenyl group.

15. (Original) The composition according to claim 12 wherein R₂, R₃, R₄, R₅ and R₆ are H.

16. (Previously presented) The composition according to claim 12 wherein R₁ is H or a C₁-C₆alkyl or cycloheteroalkyl group each optionally substituted.

17. (Original) The composition according to claim 16 having a compound of formula I selected from the group consisting of:

1-(phenylsulfonyl)-4-piperazin-1-yl-1H-indole;

1-[(2-bromophenyl)sulfonyl]-4-piperazin-1-yl-1H-indole;

1-[(6-chloroimidazo[2,1-b][1,3]thiazol-5-yl)sulfonyl]-4-piperazin-1-yl-1H-indole;


1-[(3,4-dimethoxyphenyl)sulfonyl]-4-piperazin-1-yl-1H-indole;

1-[(5-chloro-3-methyl-1-benzothien-2-yl)sulfonyl]-4-piperazin-1-yl-1H-indole;

B' 1-[(4-bromophenyl)sulfonyl]-4-piperazin-1-yl-1H-indole;
1-[(5-bromothien-2-yl)sulfonyl]-4-piperazin-1-yl-1H-indole;
1-[(4,5-dichlorothien-2-yl)sulfonyl]-4-piperazin-1-yl-1H-indole;
methyl 4-[(4-piperazin-1-yl-1H-indol-1-yl)sulfonyl]phenyl ether;
4-piperazin-1-yl-1-[[4-(trifluoromethoxy)phenyl]sulfonyl]-1H-indole;
4-(4-benzylpiperazin-1-yl)-1-(phenylsulfonyl)-1H-indole;
4-(4-benzylpiperazin-1-yl)-1-[(2-bromophenyl)sulfonyl]-1H-indole;
4-(4-benzylpiperazin-1-yl)-1-[(6-chloroimidazo[2,1-b][1,3]thiazol-5-yl)sulfonyl]-1H-indole;
4-(4-benzylpiperazin-1-yl)-1-[(3,4-dimethoxyphenyl)sulfonyl]-1H-indole;
4-[4-(3-methoxybenzyl)piperazin-1-yl]-1-(phenylsulfonyl)-1H-indole;
1-(phenylsulfonyl)-4-[4-(pyridin-4-ylmethyl)piperazin-1-yl]-1H-indole;
1-(phenylsulfonyl)-4-[4-(pyridin-3-ylmethyl)piperazin-1-yl]-1H-indole;
1-[(2-bromophenyl)sulfonyl]-4-[4-(3-methoxybenzyl)piperazin-1-yl]-1H-indole;
1-[(2-bromophenyl)sulfonyl]-4-[4-(pyridin-4-ylmethyl)piperazin-1-yl]-1H-indole;
1-[(2-bromophenyl)sulfonyl]-4-[4-(pyridin-3-ylmethyl)piperazin-1-yl]-1H-indole;
1-(phenylsulfonyl)-5-piperazin-1-yl-1H-indazole;
1-(phenylsulfonyl)-6-piperazin-1-yl-1H-indazole;
1-[(2-bromophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
1-[(4-bromophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
1-[(4-bromophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
1-[(5-bromothien-2-yl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
1-[(5-bromothien-2-yl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
1-[(4-fluorophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
1-[(4-fluorophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
methyl 4-[(5-piperazin-1-yl-1H-indazol-1-yl)sulfonyl]phenyl ether;
1-phenylsulfonyl-4-(4-propylpiperazin-1-yl)-1H-indazole;
1-phenylsulfonyl-4-piperazin-1-yl-1H-indazole;
1-phenylsulfonyl-4-(4-phenethylpiperazin-1-yl)-1H-indazole;
1-phenylsulfonyl-4-[4-(3-phenylpropyl)-piperazin-1-yl]-1H-indazole; and
the pharmaceutically acceptable salts thereof.

18. (Previously presented) A method for the preparation of a compound of formula I.

B1



(I)

wherein

A is N;

X is CR₁₁ or N;

Y is CR₇ or N with the proviso that when X is N, then Y must be CR₇;

R₁ is C₁-C₆alkylcarbonyl, C₁-C₆alkylcarbonyloxy or an C₁-C₆alkyl, C₁-C₆alkenyl, C₁-C₆alkynyl or cycloheteroalkyl group each optionally substituted;

R₂, R₃, R₄, R₅ and R₆ are each independently H, halogen, OH or an optionally substituted C₁-C₆alkyl group;

R₇ and R₁₁ are each independently H, halogen or an C₁-C₆alkyl, aryl, heteroaryl or alkoxy group each optionally substituted;

R₈ is an C₁-C₆alkyl, aryl or heteroaryl group each optionally substituted;

R₉ is H, halogen or an C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkenyl, aryl or heteroaryl group each optionally substituted;


R₁₀ is H, OH or an optionally substituted C₁-C₆alkoxy group;

m is an integer of 2;

n is 0 or an integer of 1, 2 or 3; and

--- represents a single bond or a double bond


said method which comprises reacting a compound of formula Ia



(Ia)

wherein A, X, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, m and n are as defined hereinabove for formula I with a compound R₁-Hal wherein R₁ is as defined hereinabove for formula I and Hal is Cl, Br or I.

19. (Currently amended) A compound of formula I



(I)

wherein

A is N;

X is CR₁₁ or N;

Y is CR₇ or N with the provisos that when X is N, then Y must be CR₇ and at least one of X and Y must be N;

R₁ is H, C₁-C₆alkylcarbonyl, C₁-C₆alkylcarbonyloxy or an C₁-C₆alkyl, C₁-C₆alkenyl, C₁-C₆alkynyl or cycloheteroalkyl group each optionally substituted;

R₂, R₃, R₄, R₅ and R₆ are each independently H, halogen, OH or an optionally substituted C₁-C₆alkyl group;

~~R₇~~ and R₁₁ ~~are each independently~~ is H, halogen or an C₁-C₆alkyl, aryl, heteroaryl or C₁-C₆alkoxy group each optionally substituted;

R₈ is an C₁-C₆alkyl, aryl or heteroaryl group each optionally substituted;

R₉ is H, halogen or an C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkenyl, aryl or heteroaryl group each optionally substituted;

R₁₀ is H, OH or an optionally substituted C₁-C₆alkoxy group;

m is an integer of 2;

n is 0 or an integer of 1, 2 or 3; and

--- represents a single bond or a double bond; or


a pharmaceutically acceptable salt thereof.

20. (Previously presented) The compound according to claim 19 wherein R₈ is an optionally substituted phenyl group.

21. (Previously presented) The compound according to claim 19 selected from the group consisting of:

1-(phenylsulfonyl)-5-piperazin-1-yl-1H-indazole;
1-(phenylsulfonyl)-6-piperazin-1-yl-1H-indazole;
1-[(2-bromophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
1-[(4-bromophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
1-[(4-bromophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
1-[(5-bromothien-2-yl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
1-[(5-bromothien-2-yl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
1-[(4-fluorophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
1-[(4-fluorophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
methyl 4-[(5-piperazin-1-yl-1H-indazol-1-yl)sulfonyl]phenyl ether;
1-phenylsulfonyl-4-(4-propylpiperazin-1-yl)-1H-indazole;
1-phenylsulfonyl-4-piperazin-1-yl-1H-indazole;
1-phenylsulfonyl-4-(4-phenethylpiperazin-1-yl)-1H-indazole;
1-phenylsulfonyl-4-[4-(3-phenylpropyl)-piperazin-1-yl]-1H-indazole; and
the pharmaceutically acceptable salts thereof.

22. (Currently amended) A pharmaceutical composition which comprises a pharmaceutically acceptable carrier and an effective amount of a compound of formula I.



(I)

wherein

A is N;

X is CR₁₁ or N;

Y is CR₇ or N with the proviso that when X is N, then Y must be CR₇ and at least one of X and Y must be N;

B' R₁ is H, C₁-C₆alkylcarbonyl, C₁-C₆alkylcarbonyloxy or an C₁-C₆alkyl, C₁-C₆alkenyl, C₁-C₆alkynyl or cycloheteroalkyl group each optionally substituted;

R₂, R₃, R₄, R₅ and R₆ are each independently H, halogen, OH or an optionally substituted C₁-C₆alkyl group;

~~R₇ and R₁₁ are each independently~~ is H, halogen or an C₁-C₆alkyl, aryl, heteroaryl or C₁-C₆alkoxy group each optionally substituted;

R₈ is an C₁-C₆alkyl, aryl or heteroaryl group each optionally substituted;

R₉ is H, halogen or an C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkenyl, aryl or heteroaryl group each optionally substituted;

R₁₀ is H, OH or an optionally substituted C₁-C₆alkoxy group;

m is an integer of 2;

n is O or an integer of 1, 2 or 3; and

--- represents a single bond or a double bond; or

a pharmaceutically acceptable salt thereof.

23. (Previously presented) The composition according to claim 22 having a compound of formula I selected from the group consisting of:

1-(phenylsulfonyl)-5-piperazin-1-yl-1H-indazole;

1-(phenylsulfonyl)-6-piperazin-1-yl-1H-indazole;

1-[(2-bromophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;

1-[(4-bromophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;

1-[(4-bromophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;

1-[(5-bromothiophen-2-yl)sulfonyl]-5-piperazin-1-yl-1H-indazole;

1-[(5-bromothiophen-2-yl)sulfonyl]-6-piperazin-1-yl-1H-indazole;

1-[(4-fluorophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;

1-[(4-fluorophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;

methyl 4-[(5-piperazin-1-yl-1H-indazol-1-yl)sulfonyl]phenyl ether;

1-phenylsulfonyl-4-(4-propylpiperazin-1-yl)-1H-indazole;

1-phenylsulfonyl-4-piperazin-1-yl-1H-indazole;

1-phenylsulfonyl-4-(4-phenethylpiperazin-1-yl)-1H-indazole;

1-phenylsulfonyl-4-[4-(3-phenylpropyl)-piperazin-1-yl]-1H-indazole; and

B' the pharmaceutically acceptable salts thereof.
